





# Agenda

- The Goal of Practical Feild Unification
- Dirac and Cosmic Evidence of Unification
- The Poynting Vector and Gravity
- The Quantum Vacuum
- The GEM Theory
- The Poynting Motivator Coupling to the Vacuum
- Discussion and Summary



# My Science Fiction






"Prequel" to Morningstar Pass




# Our goal: Propulsion by Manipulation of Spacetime



Scotty! More anti-matter!



#### James Clerk Maxwell



The "Great Unifier"


# MORNINGSTAR Electricity-Magnetism Unification (1861)



electricity



light




**EM Theory** 





### Einstein Sought Gravity-EM Unification



Goal: find the EM structure of spacetime- and alter it!



# Big Question:

Was Einstein partly right?:


Can one unify EM and Gravitation, both long range forces without explicit mention of the Weak and Strong Forces?

Using Quantum theory of the Vacuum?

(Which Einstein rejected)



# Dirac's BIG Discovery



$$\frac{4\pi\varepsilon_o Gm_p m_e}{e^2} \cong \frac{r_c}{R_H}$$

$$\frac{e^2}{4\pi\varepsilon_o m_o c^2} = r_c \qquad \begin{array}{c} \text{Electron classical} \\ \text{radius} \end{array}$$

 $R_H$  is the Hubble Radius

But what does it mean?




Condition of critically open cosmos:

Condition of critical optical depth

$$\Omega = \frac{8\pi G m_p n R_H^2}{3c^2} \cong 1 \qquad \cong \sigma_{Th} n R_H = \frac{8\pi n r_c^2 R_H}{3} \cong 1$$

$$\frac{e^2}{4\pi\varepsilon_o m_e c^2} = r_c$$
 Electron classical radius

Dirac Condition



(Brandenburg 1992)






# Physical Meaning of Dirac Condition?

Gravity Coupling of the particles in the universe is correlated to their EM Coupling with each other by radiation




#### Sakharov's Unification Model

Gravity as ZPF (Zero Point Fluctuation) Radiation Pressure










TWO DARK OBJECTS IN A
BRIGHT BOX
ATTRACT EACH OTHER



### Source of Sakharov's Model













Hydrogen Bomb Operates on Radiation Pressure



# Hilbert Action Principle

 General Relativity for a vacuum can be derived from spacetime energy minimization

$$W = \int \frac{R}{16\pi G} dx^4$$

$$\delta W = 0$$

Sakharov used this concept

$$R_{\mu\nu} + \frac{1}{2} g_{\mu\nu} R = 0$$



### Sakharov's Formula for G

$$G \cong \frac{c^5}{\omega_p} = \frac{c^3 r_p^2}{\hbar}$$

$$\int_0^{\hbar} \hbar \omega d\omega$$

"Big G from ZPF"

$$r_p = \sqrt{\frac{G\hbar}{c^3}}$$
 Planck Length

GRAVITY FIELDS ARE QUANTUM EM



# t'Hooft Derivation of Newton Gravitation Constant



Black Hole –pure gravity

Hawking Decay

Pure thermal EM

$$Gm_e^2 \cong (\alpha\sqrt{2})^{-1} e^{-\pi/4\alpha} (?),$$


A PHYSICAL INTERPRETATION OF GRAVITATIONAL INSTANTONS

G. 't HOOFT Nuclear Physics B315 (1989) 517-527 North-Holland, Amsterdam



# Common Concept among ideas

Gravity vector is connected to the EM Poynting vector



$$\vec{S} = \frac{\vec{E}x\vec{B}}{\mu_o}$$



## Kaluza-Klein Theory

• Add a compact 5<sup>th</sup> dimension to the Hilbert Action principle

Obtain General Relativity Plus Maxwell's

equations






Theodor Kaluza and Oskar Klein



## Kaluza-Klein Theory (1929)





## Kaluza Klein Action principle

$$W = \int K dx^{4}$$

$$\delta W = 0$$

$$K = \frac{R}{16\pi G} - \frac{F_{\mu\gamma}F^{\mu\gamma}}{4}$$





# Introduction To GEM Theory

- GEM (Gravity-Electro-Magnetism)or (Grandis et Medianis) ) the "great and the middle" scales
- GEM is an alloy of Sahkarov -ZPF (Zero Point Fluctuation), and Kaluza Klein (5<sup>th</sup> Dimensional) approachs to EM-gravity unification



# Planck's Constant and Quantum mechanics

• Planck scale is length where spacetime is a foam of quantum black holes appearing and annihilating each other because of Heisenberg Uncertainty:

$$r_{\rm P} = 1.6 \times 10^{-33} \text{ cm}$$
 $r_{\rm P} = \sqrt{\frac{G\hbar}{c^3}}$ 
 $r_{\rm P} = \sqrt{\frac{G\hbar}{c^3}}$ 

• EM turbulence caused by this Hiesenberg Uncertainty gives rise to ZPF of EM fields





### The Cosmos as it is

- Cosmos is 99.9% hydrogen (protons and electrons)
- It has a heavy particle, a proton, and lighter one, electron (yin and yang?)
- It has two long range forces: weak one-gravity and strong one-EM  $\frac{Gm_pm_e}{e^2} = 4x10^{-40}$
- Gravity satisfies equivalence, EM doesn't



#### The Mesoscale

- Mesoscale is the scale of ordinary subatomic particles and their scale lengths.
  - masses:  $\sim m_p$ ,  $m_e$ ,
  - lengths:  $\sim r_c = e^2/(m_e c^2)$
- Mesoscale is between Planck scale and Cosmic scale in length
- Mesoscale below Planck scale in mass



# The GEM Theory (1986)

- Two Postulates:
  - Gravity is an array of Poynting vectors (ExB drifts)
  - Both EM- gravity and protons and electrons appear from the Planck scale with the appearance of a 5<sup>th</sup> dimensional hidden dimension

**GEM Theory combines Sakharov and Kaluza-Klein Theories** 



# A Model for Gravity fields in terms of E and B fields

$$V_D = \frac{(\vec{E}x\vec{B})}{B^2} = \frac{\mu_o \vec{S}}{B^2} \qquad \blacksquare \qquad E \qquad \blacksquare$$


$$g = \frac{\dot{E}xB_o c}{B_o^2} \qquad \qquad U_d \qquad \blacksquare$$

ExB drift effects all particles regardless of mass or charge, satisfies equivalence principle



Uniform motion


# ExB drift Gravity Cont.



Acceleration



# ExB drift Gravity Cont.



We have model of gravity field as EM for "free particles"

Simulation using EM PIC code



### Vacuum mass

$$g_{\alpha\beta} = \frac{4F_{\alpha}^{\gamma}F_{\gamma\beta}}{F_{\mu\gamma}F^{\mu\gamma}}$$

$$T_{\alpha\beta} = F_{\alpha}^{\gamma} F_{\gamma\beta} - g_{\alpha\beta} \frac{F_{\mu\gamma} F^{\mu\gamma}}{4} = 0$$

The Vacuum Weighs Nothing!!!!!!

But is full of energy!



### **GEM Metric Tensor**

$$g_{\alpha\beta} = \frac{4F_{\alpha}^{\gamma}F_{\gamma\beta}}{F_{\mu\gamma}F^{\mu\gamma}} = \left[B^{2} - E^{2}\right]^{-1} \begin{pmatrix} -E^{2} & S_{x} & S_{y} & S_{z} \\ S_{x} & E_{x}^{2} - B_{y}^{2} - B_{z}^{2} & & & \\ S_{y} & E_{y}^{2} - B_{x}^{2} - B_{z}^{2} & & & \\ S_{z} & & E_{z}^{2} - B_{y}^{2} - B_{x}^{2} \end{pmatrix}$$

$$S = ExBc$$



### Size of Hidden Dimension

$$\phi_o(\eta) = \ln(\ln(r_c(\xi)/r_P))$$

$$\left(\frac{m_p}{m_e}\right)^{1/2} = \ln\left(\frac{r_o}{r_p}\right) = 42.8503...$$

The postulate that the separation of proton-electron pair into distinct particles is correlated to the separation of gravity and EM forces- leads to relation of mass ratio to ratio of strengths



# Vacuum Bernoulli Equation

Kaluza Klien Action

$$K = \frac{R}{16\pi G} - \frac{F_{\mu\gamma}F^{\mu\gamma}}{4} \Rightarrow \frac{g^2}{2\pi G} - \frac{S^2}{U_o}$$

$$U_o = \frac{(B^2 - E^2)}{8\pi}$$

$$\frac{g^2}{2\pi G} - \frac{S^2}{U} = K \quad Vacuum \quad Bernoulli \quad Equation$$

Direct coupling between EM and gravity fields



### **GEM Results**

#1 
$$\frac{g^2}{2\pi G} - \frac{S^2}{U} = K$$
 Vacuum Bernoulli Equation

Direct coupling between EM and gravity fields

#2 
$$\ln \left[ \frac{r_o}{r_P} \right] = \left[ \frac{m_p}{m_e} \right]^{1/2} = 42.8503...$$

Mass formula leads to formula for "big G"



### Theoretical formula for G

$$G = \frac{e^2}{4\pi\varepsilon_o m_p m_e} \alpha \left[\frac{r_p}{r_o}\right]^2$$



$$G = \frac{e^2}{4\pi\varepsilon_o m_p m_e} \alpha \exp(-2\left[\frac{m_p}{m_e}\right]^{1/2}) = 6.668 \times 10^{-11} N - m^2 / kg^2$$

Inversion of log formula leads to formula for "big G" (in MKS)

Accepted value:

 $G = 6.673x10^{-11}m^3kg^{-1}\sec^{-2}\pm 1.5$  parts per thousand (CODATA)



# Theoretical Versus Accepted



$$G = 6.673x10^{-11}m^3kg^{-1}\sec^{-2}\pm 1.5$$
 parts per thousand (CODATA)

$$\frac{G_{\text{exp}} - G_{\text{theory}}}{G_{\text{exp}}} = \frac{6.673 \times 10^{-11} - 6.6680 \times 10^{-11}}{6.6.73 \times 10^{-11}} = .0007439 = 0.7 \times 10^{-3} error$$

Formula for G is within experimental error to accepted value



#### Vacuum Hall Effect

GEM theory predicts Poynting flows can couple to vacuum "Virtual Plasma"

Momentum conservation happens through gravity coupling of vaccum to nearby large masses



# Parallel Universes Theory (2014)

New formulation of Quantum Mechanics

Vacuum consists of multiple-coexisting 'spaces' some just after their own "Big Bang" and *thus full of plasma* 



Theory allows propulsion Poynting fields to tunnel to parallel universe full of plasma and reaction fields to tunnel back allowing coupling to "vacuum"



### **SUMMARY**

- GEM theory is combination of Sakharov and Kaluza-Klein
- GEM theory is covariant
  - Metric tensor is normalized EM stress tensor
  - Gravity vector associated with Poynting Vector
  - ZPF mass vanishes but fields do not
- Accurate formula for is found

  - "Vacuum Hall Effect" is predicted in GEM theory making "propellant-less" propulsion possible.






### There is no "Box"







I told you so!